



STATE RESEARCH CENTER OF RUSSIA
INSTITUTE FOR HIGH ENERGY PHYSICS

IHEP 98-62

D.V. Amelin, E.B. Berdnikov, S.I. Bityukov,
G.V. Borisov, V.A. Dorofeev, R.I. Dzhelyadin,
A.V. Ekimov, Yu.P. Gouz, I.A. Kachaev,
A.N. Karyukhin, Yu.A. Khokhlov, A.K. Konoplyannikov,
V.F. Konstantinov, S.V. Kopikov, M.E. Kostrikov,
V.V. Kostyukhin, S.A. Likhoded, V.D. Matveyev,
A.P. Ostankov, D.I. Ryabchikov, A.A. Solodkov,
O.V. Solovianov, E.A. Starchenko, N.K. Vishnevsky,
E.V. Vlassov, A.M. Zaitsev
IHEP, Protvino
G.G. Sekhniadze, E.G. Tskhadadze,
IPh, Tbilisi

**STUDY OF $\eta\pi^-$ PRODUCTION BY PIONS IN THE
COULOMB FIELD**

Submitted to *Yad. Fiz.*

Protvino 1998

Abstract

Amelin D.V. et al. Study of $\eta\pi^-$ production by pions in the Coulomb field: IHEP Preprint 98-62. – Protvino, 1998. – p. 8, figs. 5, refs.: 7.

The production of $\eta\pi^-$ system at low $M_{\eta\pi^-}$ by the π^- beam in the Coulomb field of Be nuclei was studied. The cross section measurements of the reaction $\pi^-Be \rightarrow \eta\pi^-Be$ were compared with the experimental data on the decay $\eta \rightarrow \pi^+\pi^-\gamma$ and the theoretical predictions.

Аннотация

Д.В. Амелин и др. Исследование образования системы $\eta\pi^-$ пионами в кулоновском поле: Препринт ИФВЭ 98-62. – Протвино, 1998. – 8 с., 5 рис., библиогр.: 7.

На установке ВЕС (ИФВЭ, Протвино) исследовано образование системы $\eta\pi^-$ с малыми массами пучком пионов в кулоновском поле ядра бериллия. Результаты измерения сечения реакции $\pi^-Be \rightarrow \eta\pi^-Be$ сравниваются с экспериментальными данными о ширине распада $\eta \rightarrow \pi^+\pi^-\gamma$ и с теоретическими моделями.

1. Introduction

We present here the study of the $\eta\pi^-$ system production at low mass ($M_{\eta\pi^-} < 1.18 \text{ GeV}/c^2$) in the Coulomb field of Be nuclei

$$\pi^- Be \rightarrow \eta\pi^- Be. \quad (1)$$

The subject of our study is the process

$$\pi^- \gamma \rightarrow \eta\pi^-. \quad (2)$$

Its amplitude at low $M_{\eta\pi^-}$ can be expressed in the following form:

$$M_{\eta\pi\pi\gamma} = -i\epsilon_{\mu\nu\rho\sigma} A^\mu p_i^\nu p_\pi^\rho p_\eta^\sigma F_{\eta\pi\pi\gamma}(s, t, u) \quad (3)$$

(notations are given in Fig.1, $s = (p_\pi + p_\eta)^2$, $t = (p_i - p_\pi)^2$, $u = (p_i - p_\eta)^2$).




Fig. 1. Reaction (1).

In the chiral limit the amplitude of process (2) is determined by the box anomaly, and $F_{\eta\pi\pi\gamma}(0,0,0)$ can be expressed as follows [1,2]:

$$F_{\eta\pi\pi\gamma}(0,0,0) = \frac{e}{4\pi^2 f_\pi^3} \left(\frac{f_\pi \cos\theta_p}{f_8 \sqrt{3}} - \frac{f_\pi}{f_0} \sqrt{\frac{2}{3}} \sin\theta_p \right). \quad (4)$$

Here $f_{\pi,0,8}$ are the pion, singlet and octet decay constants, θ_p is the singlet-octet mixing angle for pseudoscalars.

The value of $F_{\eta\pi\pi\gamma}$ at the chiral limit as well as its dependence on the kinematical variables are of interest for the theoretical analysis.

A model predicting the dependence of $F_{\eta\pi\pi\gamma}(s,t,u)$ on the kinematical variables was described in [2]; following this model, the authors performed the fit of experimental data on radiative decays of light pseudoscalar mesons

$$\eta, \eta' \rightarrow \pi^+ \pi^- \gamma, \quad (5)$$

$$\eta, \eta' \rightarrow \gamma \gamma \quad (6)$$

and determined f_0 , f_8 and θ_p . For $F_{\eta\pi\pi\gamma}$ this fit yields

$$F_{\eta\pi\pi\gamma}(0,0,0) = 6.5 \pm 0.3 \text{ GeV}^{-3}. \quad (7)$$

Decays (5) proceed at positive $t = M_{\pi\pi}^2$. Study of reaction (2) gives data on $F_{\eta\pi\pi\gamma}(s,t,u)$ in other region of kinematical variables, at negative t . Here we compare the experimentally measured cross section of reaction (1) at low $M_{\eta\pi^-}$ with the predictions from [2].

2. Main features of the $\eta\pi^-$ system production in the Coulomb field of nuclei

The cross section of the $\eta\pi^-$ system production in the Coulomb field of nuclei is given by expression [3]

$$\frac{d\sigma}{dsdq^2} = \frac{Z^2 \alpha}{\pi} \frac{|q^2 - q_{min}^2|}{q^4} \frac{1}{s - m_\pi^2} \frac{p_\eta^3 p_i}{48\pi} F_{\eta\pi\pi\gamma}^2 G^2(q^2). \quad (8)$$

Here Z is the charge of nucleus, $G(q^2)$ is its electromagnetic form-factor, p_i and p_η are the absolute values of momenta of an incident pion and outgoing η -meson in the c.m. system of produced η and π^- .

The differential cross section of this reaction is strongly peaking at low q^2 . This feature facilitates the event selection; on the other hand, it leads to the fast drop of the cross section at higher masses of produced $\eta\pi^-$ due to the cut on $|q_{min}^2| \simeq \frac{(s - m_\pi^2)^2}{4E_{beam}^2}$.

In case of the $\eta\pi^-$ production at low mass in the Coulomb field, the P -wave is predominantly produced¹ (S -wave corresponds to 0-0 transition and is suppressed $\propto q^2/m_\rho^2$,

¹Note that the P -wave in the $\eta\pi^-$ system has an exotic set of quantum numbers, $I^C J^P = 1^- 1^-$.

and higher waves are suppressed by the barrier factor p^{2l}). However, higher waves appear in the amplitude (3) due to the dependence of $F_{\eta\pi\pi\gamma}$ on kinematical variables, and their fraction increases with $M_{\eta\pi^-}$.

The background for the process of production in the Coulomb field is the process of hadronic production of the $\eta\pi^-$ system. In the hadronic production of $\eta\pi^-$ at high energies the amplitudes with a positive exchange naturality with the projection of 1 onto the Gottfried-Jackson axis are dominating [4]. At low $|q^2|$ these are mainly diffractive processes, which proceed coherently on nuclei. They have a characteristic q^2 distribution in the form

$$\frac{d\sigma}{dq^2} \sim |q^2 - q_{min}^2| e^{-b|q^2|} \quad (9)$$

with the slope parameter $b \simeq 50 \text{ GeV}^{-2}$ and the distribution on an azimuthal angle in the Gottfried-Jackson frame (Treiman-Yang angle)

$$\frac{dN}{d\varphi_{TY}} \sim \sin^2 \varphi_{TY}. \quad (10)$$

The $\eta\pi^-$ can also be produced in processes with negative exchange naturality (e.g. by b_1 -trajectory). Such exchanges, however, are suppressed in scattering on nuclei at high energies, which can be illustrated by the absence of a signal from $a_0(980)$ -meson in the $\eta\pi^-$ effective mass spectrum [4]. At low $M_{\eta\pi^-}$ for the contribution of such exchanges one can expect zero projection onto the Gottfried-Jackson axis (uniform distribution on Treiman-Yang angle) and the broad q^2 -distribution of the type

$$\frac{d\sigma}{dq^2} \sim e^{-b|q^2|}. \quad (11)$$

with the slope parameter $b \simeq 7 \text{ GeV}^{-2}$. As the contribution of negative naturality exchanges is expected to be small, such approximate description is sufficient for our studies.

Apart from hadronic production of $\eta\pi^-$, there exists an instrumental background, which consists of events with different final states, where one or more slow particles have not been detected. For this background one can expect the broad q^2 -distribution, which can also be parametrized in the form of $d\sigma/dq^2 \sim e^{-b|q^2|}$.

3. The VES setup

The experiment is performed at the VES setup (IHEP, Protvino) with π^- beam at the momentum of $37 \text{ GeV}/c$. The setup is a large aperture magnetic spectrometer which includes a system of proportional and drift chambers and a lead-glass electromagnetic calorimeter. The target was beryllium ($l = 4 \text{ cm}$). The trigger conditions required the presence of two or more charged particles in the forward hemisphere and the absence of hard charged particles in the backward hemisphere. The process under study (1) was detected with η -meson decaying into $\pi^+\pi^-\pi^0$. In the region of $M_{\eta\pi^-} = 1 \text{ GeV}/c^2$, the setup resolution on the effective mass of $\eta\pi^-$ is $\sigma(M_{\eta\pi^-}) \simeq 15 \text{ MeV}/c^2$ and on the transverse momentum $\sigma(p_t) \simeq 17 \text{ MeV}/c$.

4. Event selection, results

The main event selection criteria were the following:

- the event contains three tracks of charged particles with a total charge of -1 and two photons in the final state;
- the total energy of the final state lies within the interval of $36 < E_{tot} < 39$ GeV;
- the charged particles are not identified as electrons;
- the effective mass of two photons lies within the interval of the π^0 mass: $105 < m_{\gamma\gamma} < 165$ MeV/ c^2 ;
- the effective mass of one of $\pi^+\pi^-\pi^0$ subsystems lies in the interval of the η -meson mass: $531 < m_{\pi^+\pi^-\pi^0} < 567$ MeV/ c^2 . The constraint $m_{\gamma\gamma} = m_{\pi^0}$ was applied for the calculation of $m_{\pi^+\pi^-\pi^0}$.

Fig.2 shows $m_{\gamma\gamma}$ (a) and $m_{\pi^+\pi^-\pi^0}$ (b) spectra. Signals from π^0 and η mesons are clearly seen against a smooth background.




Fig. 2. Effective mass spectra of $\gamma\gamma$ (a) and $\pi^+\pi^-\pi^0$ (b). The cuts used for the event selection are shown by arrows.

Fig.3 shows the $M_{\eta\pi^-}$ spectrum. It is dominated by the $a_2^-(1320)$ -meson signal. The arrow shows the cut used for the selection of events near the threshold: $M_{\eta\pi^-} < 1.18$ GeV/ c^2 . The background for this spectrum was evaluated and subtracted using the events from reference intervals on $m_{\pi^+\pi^-\pi^0}$ around the η -meson peak, with $503 < m_{\pi^+\pi^-\pi^0} < 521$ MeV/ c^2 or $577 < m_{\pi^+\pi^-\pi^0} < 595$ MeV/ c^2 . For all other spectra the method of bin filtering was used, i.e. the number of events in each bin was determined as the number of η -mesons obtained by fitting the $m_{\pi^+\pi^-\pi^0}$ spectrum for this bin.

Fig.4 shows the $|q^2 - q_{min}^2|$ distribution for the selected events. It has a characteristic Coulomb peak at low $|q^2 - q_{min}^2|$, as well as a broad structure corresponding to the hadronic production of $\eta\pi^-$. The result of the fit by the superposition of q^2 -distributions for the Coulomb (8) and hadronic (9) production of $\eta\pi^-$, convoluted with the experimental resolution of the setup and the instrumental background is shown by the dashed line in Fig.4.




Fig. 3. The effective mass spectrum of $\eta\pi^-$. The arrow shows the cut used for the event selection ($M_{\eta\pi^-} < 1.18 \text{ GeV}/c^2$).




Fig. 4. The $|q^2 - q_{min}^2|$ distribution for events with $m_{\eta\pi^-} < 1.18 \text{ GeV}/c^2$. The dashed line is the result of the fit (see text); the dotted line shows the contribution of hadronic production of $\eta\pi^-$ and background.

The parameters of the instrumental background were determined by the analysis of distributions on the total energy and Treiman-Yang angle for the events with $0.01 < |q^2 - q_{min}^2| < 0.1 \text{ GeV}^2$ and $M_{\eta\pi^-} < 1.18 \text{ GeV}/c^2$ (Fig. 5a,b; in the first case the aforementioned selection on the total energy was not applied). In the total energy distribution one can see the peak corresponding to fully reconstructed events. The background to the left of the peak, with $34 < E_{tot} < 35.5 \text{ GeV}$, has the uniform distribution on the Treiman-Yang angle

and q^2 distribution with slope parameter $b \simeq 7 \Gamma \text{eB}^{-2}$. The number of background events was determined by fitting the distribution on Treiman-Yang angle by the superposition of (10) and the constant. It is worth noticing that similar distributions on φ_{TY} and $|q^2|$ are also expected for $\eta\pi^-$ production in isospin exchanges with negative naturality, which means that the number of background events obtained by fitting the Treiman-Yang angle distribution also contains a possible contribution from such exchanges.




Fig. 5. Distributions on total energy (a) and Treiman-Yang angle (b) for the $\eta\pi^-$ events with $M_{\eta\pi^-} < 1.18 \text{ GeV}/c^2$, $0.01 < |q^2 - q_{min}^2| < 0.1 \text{ GeV}^2$.

The number of events in the Coulomb peak, in the range of $|q^2 - q_{min}^2| < 0.09 \text{ GeV}^2$, is

$$N_{coul} = 109 \pm 23. \quad (12)$$

Free parameters of the fit to the q^2 -distribution (Fig. 4) were the number of events of the Coulomb production of $\eta\pi^-$ (8) and the number of events of its hadronic production. The interference of Coulomb and hadronic amplitudes was taken to be negligible, because the former is real, while the latter, at high energies, is almost imaginary. For the slope parameter b in (9) the value of $b = 50 \text{ GeV}^{-2}$ was used, which corresponds to the diffractive production on Be nuclei. The variation of this value by $\pm 10\%$ does not change the result. The number of events of the instrumental background was taken equal to that found from fitting the distribution on Treiman-Yang angle. The error in the determination of free parameters accounts for the influence of variation in the background events number within its standard deviation.

In order to find the cross section, we used the results of [5]², where the differential cross section of the reaction

$$\pi^- Be \rightarrow \pi^+ \pi^- \pi^- Be \quad (13)$$

²Authors express their thanks to V.V. Ezhela and Yu.I. Ivanshin for the help in extracting the necessary data from [5].

was measured at 40 GeV/c, and also the results of the VES experiment on reactions

$$\begin{aligned}\pi^- Be &\rightarrow \pi^+ \pi^- \pi^- Be \\ \pi^- Be &\rightarrow a_2^-(\pi^+ \pi^- \pi^-) Be \\ \pi^- Be &\rightarrow a_2^-(\eta(\pi^+ \pi^- \pi^0) \pi^-) Be.\end{aligned}$$

These data allow us to determine the cross section:

$$\sigma_{coul} = 145 \pm 34 \text{ nb.} \quad (14)$$

When calculating the error, we took into account the error in the measurements of [5] (5%) and the systematic error (10%) which corresponds to the uncertainty in the calculation of the VES setup efficiency.

Besides the process under study (2), the Coulomb production of a_2^- -meson decaying into $\eta\pi^-$ also contributes to this cross section. Assuming a positive interference between the amplitude (3) and that of the Coulomb production of $\eta\pi^-$ via a_2^- -meson, one can calculate expected cross section of the Coulomb production in the same range of $M_{\eta\pi^-}$ and q^2 , using the experimentally measured [6] width of the decay $a_2^-(1320) \rightarrow \pi^- \gamma$ and the parametrization of $F_{\eta\pi\pi\gamma}$ dependence on s, t, u from [2]:

$$\sigma_{\eta\pi}^{th} = 119 \pm 13 \text{ nb,} \quad (15)$$

(the quoted error corresponds to the errors of experimental measurements of widths of $a_2^-(1320) \rightarrow \pi\gamma$ and $\eta \rightarrow \pi^+ \pi^- \gamma$ decays). This value is in good agreement with the experimentally measured cross section (14) and confirms a strong dependence of $F_{\eta\pi\pi\gamma}$ on t .

For the description of the dependence of $F_{\eta\pi\pi\gamma}$ on kinematical variables in various theoretical models [1,2], the two substantial parameters are used — the value of $F_{\eta\pi\pi\gamma}$ in the chiral limit and the mass of vector meson in the $\pi\pi$ channel (ρ -meson). Using the cross section measured in our experiment (14) and the width of the decay $\eta \rightarrow \pi^+ \pi^- \gamma$, we calculated $F_{\eta\pi\pi\gamma}(0, 0, 0)$ and the effective mass of vector meson \tilde{m}_ρ within the framework of model [2]. We obtained the values

$$F_{\eta\pi\pi\gamma}(0, 0, 0) = 6.9 \pm 0.7 \text{ GeV}^{-3}, \quad \tilde{m}_\rho = 900 \pm 120 \text{ MeV}/c^2 \quad (16)$$

which agree with (7) and the ρ -meson mass from [7] within one standard deviation respectively.

Conclusions

The cross section of the $\eta\pi^-$ production at low mass in the Coulomb field of beryllium nuclei at the beam momentum $p_{beam} = 37 \text{ GeV}/c$, $M_{\eta\pi^-} < 1.18 \text{ GeV}/c^2$ and $|q^2| < 0.09 \text{ GeV}^2$ was measured

$$\sigma_{coul} = 145 \pm 34 \text{ nb.} \quad (17)$$

This value is in a good agreement with theoretical predictions [2,6].

This work was supported by Russian Foundation for Basic Research (grant 98-02-16392) and INTAS-RFBR (grant 95-0267).

References

- [1] P. Ko, T.N. Truong, Phys. Rev. D42 (1990), 2419–2422.
- [2] E.P. Venugopal and Barry R. Holstein, Phys. Rev. D57. (1998) 4397.
- [3] A. Halprin, C.M. Anderson and H. Primakoff, Phys. Rev. 152 (1966) 1295.
- [4] G.M.Beladidze et al, Phys. Lett. B313 (1993) 276–282.
- [5] G. Bellini et al, Nucl. Phys. B199 (1982) 1–26.
- [6] S. Cihangir et al, Phys. Lett. B117 (1982) 119–122.
- [7] Particle Data Group, Review of Particle Properties, Phys. Rev. D54 (1996).

Received September 24, 1998

Д.В.Амелин.

Исследование образования системы $\eta\pi^-$ пионами в кулоновском поле.

Оригинал-макет подготовлен с помощью системы \LaTeX .

Редактор Е.Н.Горина.

Технический редактор Н.В.Орлова.

Подписано к печати 08.10.1998 г. Формат $60 \times 84/8$. Офсетная печать.

Печ.л. 1. Уч.-изд.л. 0.76. Тираж 100. Заказ 310. Индекс 3649.

ЛР №020498 17.04.97.

ГНЦ РФ Институт физики высоких энергий
142284, Протвино Московской обл.

